



**Manuscript version: Author's Accepted Manuscript**

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

**Persistent WRAP URL:**

<http://wrap.warwick.ac.uk/118019>

**How to cite:**

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk).

Fluctuation and Noise Letters  
 © World Scientific Publishing Company

## Introduction to the UPoN-2018 Special Issue of *FNL*

I.A. Khovanov<sup>1\*</sup> and P.V.E. McClintock<sup>2†</sup>

<sup>1</sup>*School of Engineering, University of Warwick, Coventry CV4 7AL, UK*

<sup>2</sup>*Department of Physics, Lancaster University, Lancaster LA1 4YB, UK.*

Received (received date)

Revised (revised date)

Accepted (Day Month Year)

Noise is universal and multi-faceted. In everyday life, the aspects we encounter tend to be annoying, for example, loud conversations when we wish to enjoy a quiet cup of tea, or students chatting during a lecture. But acoustic noise can sometimes be useful e.g. when it indicates that the kettle is boiling, as an essential prelude to the tea, or when it helps to awaken somnolent students at the end of the lecture thereby eliminating the mutual embarrassment of needing to awaken them.

Similarly in science, noise as random fluctuations appears in different guises with a great diversity of origins. However, there is one feature in common to all manifestations of noise: it appears as the outcome of interactions between a huge number of components, such as charges in electronic devices, or atoms and molecules in proteins, or oscillators in an ensemble, or in an urban environment arising from the daily activities of people. This common feature defines the main challenges in the characterization and understanding of noise. Consequently, the open unsolved problems of noise are of wide applicability and naturally attract the attention of researchers from many different fields including some of the brightest and best from every generation of scientists.

The recent 8<sup>th</sup> *International Conference on Unsolved Problems of Noise* (UPoN–2018), which was organised by Janusz Smulko at the Gdańsk University of Technology, Poland, during 9–13 July 2018, provided a platform for the research community to formulate, discuss and suggest ways tackling the unsolved problems of noise. The meeting was dedicated to Michael F. Shlesinger to celebrate his 70<sup>th</sup> birthday, and in acknowledgement of his huge contributions to the subject area over the years, both scientifically and through his prescient provision of ONR funding to support research projects in stochastic nonlinear dynamics in the era before the potential

\*Email: i.khovanov@warwick.ac.uk

†Email: p.v.e.mcclintock@lancaster.ac.uk

technological importance of the subject was generally appreciated.

The UPoN conferences were initiated by Laszlo Kish, founder and Honorary Editor of *Fluctuation and Noise Letters*. The first UPoN meeting was in Szeged, Hungary, in 1996. Since then, there have been successor UPoNs in Adelaide (1999), Bethesda (2002), Gallipoli (2005), Lyon (2008), Kolkata (2012) and Barcelona (2015), leading up to UPoN-2018 in Gdańsk. In common with previous UPoNs, the meeting in Gdańsk included talks, presentations and discussions covering a large variety of noise topics. Following the UPoN tradition, the talks and discussions were often provocative, controversial, and very stimulating. This Special Issue brings together a representative selection of the topics discussed (except for those on noise in metrology, which are being published in *Metrology and Measurement Systems*).

A number of papers are devoted to noise in electronic devices. Methods for noise characterization are presented by Lentka and Smulko [1] and Ciofi *et al* [2]. Such methods are key to the comprehensive description of the stochastic output of devices. Experimental features of noise in electron transport in manganite single crystals are discussed by Przybytek and Jung [3], raising many open questions in search of theoretical explanations. Tretjak *et al* [4] have performed a comprehensive study of the noise properties of hybrid polymer composites, which are important components in modern bioelectronic devices. These composites include multi-walled carbon nanotubes and onion-like carbon particles, the presence of which defines strong temperature dependences in resistivity noise and provides an opportunity to control the noise characteristics. Experimental manifestations together with contemporary theoretical understanding of macroscopic telegraph noise in strongly correlated mesoscopic systems are discussed by Jung [5]: they describe superconductors and low Ca-doped manganites with colossal magnetoresistive as examples of systems that exhibiting similar macroscopic noise, but whose underlying noise-generating mechanisms are profoundly different.

Application of stochastic descriptions to complex deterministic systems are presented in the papers of Goldobin [6] and Dubkov *et al* [7]. In the former case, cumulant analysis – one of the basic tools in the description of noise – is applied to characterize phase dynamics. This approach allows the author to develop the theoretical framework for a deep characterization of the complex phenomena observed in a large ensemble of coupled oscillators. The latter paper presents a stochastic description of low-dimensional chaotic dynamics. The authors conclude that billiard-like particles interacting with harmonically oscillating boundaries can be considered as particles moving under the influence of white Gaussian noise whose intensity depends on the velocity of the boundary oscillations and the mean free path of the particles. This claim is supported by the results of numerical simulations.

Several papers tackle noise in biological problems. Shinagawa and Sasaki [8] provide a theoretical description of torque-enhanced diffusion in molecular motors in the presence of noise. The results of numerical calculations by the authors suggest a novel way of extracting certain properties of F<sub>1</sub>-ATPase from the enhanced

diffusion. Cugliandolo *et al* [9] consider the molecular motor as an example of an active Brownian particle, and present an approach for extracting an effective temperature based on deviations from the fluctuation dissipation theorem, illustrating the method through simulations of Langevin equations. Siódmiak and Beldowski [10] study noise spectra in hydrogen bonding energy for different temporal scales, thereby illustrating the role of phospholipids in synovial fluid. The spectra showed power-law-scaling with a large sensitivity to the presence of phospholipids that can be used for the characterisation of the chemical content of synovial fluid. Barabash *et al* [11] discuss a theoretical framework for describing the experimentally measured currents in narrow biological ion channels. Validated by molecular dynamics and Brownian dynamics simulations, it is based on a quasiparticle representation of multi-ion interactions and thus provides a direct link to approaches widely used in condensed matter physics. A non-equilibrium and self-consistent multi-species kinetic model presented by Gibby *et al* [12] aims to describe the anomalous mole fraction effect which is observed in ion channels under varying concentrations of two different ionic species. The model developed explains very well the current in a single NaChBac sodium channel and is able to describe the influence of concentration difference in sodium and potassium ions on measured currents. Vadai *et al* [13] apply spectral analysis of  $1/f$ -type noise to characterize scaling laws in human mobility. Atypically, a  $1/f$  scaling is observed for the high frequency part of the spectrum, opening up the challenging task of developing a theoretical description of this unusual scaling.

We hope that readers will enjoy perusing this Special Issue, which we believe conveys some flavour of the remarkable diversity of unsolved, and partly-solved, topical noise problems tackled during UPoN-2018.

## References

- [1] L. Lentka and J. Smulko, A simplified method of trend removal to determine noise observed during a supercapacitor's discharging, *Fluct. Noise Lett.* **18** (2019) 1940001.
- [2] C. Ciofi, G. Scandurra, and G. Giusi, QLSA: A software library for spectral estimation in low frequency noise measurements applications, *Fluct. Noise Lett.* **18** (2019) 1940004.
- [3] J. Przybytek and G. Jung, Noise features of metastable resistivity states in  $\text{La}_{0.86}\text{Ca}_{0.14}\text{MnO}_3$  manganite single crystal, *Fluct. Noise Lett.* **18** (2019) 1940011.
- [4] M. Tretjak, S. Pralgauskaitė, J. Matukas, J. Macutkevič, J. Banys, and O. Shenderova, Low frequency noise and resistivity characteristics of hybrid composites with onion-like carbon and multi-walled carbon nanotubes, *Fluct. Noise Lett.* **18** (2019) 1940009.
- [5] G. Jung, Macroscopic random telegraph noise, *Fluct. Noise Lett.* **18** (2019) 1940003.
- [6] D.S. Goldobin, Relationships between the distribution of Watanabe–Strogatz variables and circular cumulants for ensembles of phase elements, *Fluct. Noise Lett.* **18** (2019) 1940002.
- [7] A. Dubkov, A. Krasnova, and O. Chichigina, Influence of harmonic perturbation on speed of billiard particle as combination of deterministic acceleration and white noise, *Fluct. Noise Lett.* **18** (2019) ??.

- [8] R. Shinagawa and K. Sasaki, Enhanced diffusion in a model of molecular motors with potential switching, *Fluct. Noise Lett.* **18** (2019) 1940005.
- [9] L.F. Cugliandolo, G. Gonnella, and I. Petrelli, Effective temperature in active Brownian particles, *Fluct. Noise Lett.* **18** (2019) 1940008.
- [10] J. Siódmiak and P. Beldowski, Hyaluronic acid dynamics and its interaction with synovial fluid components as a source of the color noise, *Fluct. Noise Lett.* **18** (2019) ??.
- [11] M.L. Barabash, W.A.T. Gibby, C. Guardiani, D.G. Luchinsky, and P.V.E. McClintock, From the potential of the mean force to the quasiparticle's effective potential in narrow ion channels, *Fluct. Noise Lett.* **18** (2019) 1940006.
- [12] W.A.T. Gibby, M. L. Barabash, C. Guardiani, D.G. Luchinsky, O.A. Fedorenko, S.K. Roberts, and P.V.E. McClintock, Theory and experiments on multi-ion permeation and selectivity in the NaChBac ion channel, *Fluct. Noise Lett.* **18** (2019) 1940007.
- [13] G. Vadai, A. Antal, and Z. Gingl, Spectral analysis of fluctuations in humans' daily motion using location data, *Fluct. Noise Lett.* **18** (2019) 1940010.